

Long-term Effects of Neonatal Exposure to Hydroxylated Polychlorinated Biphenyls in the BALB/cCrgl Mouse

**Jeanelle M. Martinez, L. Clifton Stephens,
and Lovell A. Jones**

**doi:10.1289/ehp.7735 (available at <http://dx.doi.org/>)
Online 20 April 2005**



**Long-term Effects of Neonatal Exposure to Hydroxylated Polychlorinated
Biphenyls in the BALB/cCrgl Mouse**

Jeanelle M. Martinez,^{1,3} L. Clifton Stephens,² and Lovell A. Jones¹

Departments of Gynecologic Oncology¹, Veterinary Medicine and Surgery²,

The University of Texas, M. D. Anderson Cancer Center,

Houston, Texas 77030

³Present address and to whom correspondence should be addressed: National Institute of Environmental Health Sciences, PO Box 12233, MD C4-05, Research Triangle Park, NC 27709. Telephone: 919-541-3466. Fax: 919-541-4702. Email: martine2@niehs.nih.gov

Running Title: DEVELOPMENTAL CARCINOGENIC EFFECTS OF OH-PCBS

Key Words: hydroxylated polychlorinated biphenyls; BALB/cCrgl mouse; female reproduction; estrogenicity; tumorigenicity.

ACKNOWLEDGEMENTS: This research was funded by The Laboratory of Experimental Gynecology in the Department of Gynecologic Oncology, UTMDACC. The Center of BioEnvironmental Research, Tulane/Xavier University and NIGMS 16652. We sincerely thank Dr. Joseph K. Haseman and Dr. Grace E. Kissling for their help in all of the statistical analysis. We thank Dr. Gary Boorman for his confirmation of tumor pathology. We thank Retha Newbold and Dr. Thomas Eling for their review of this manuscript.

ABBREVIATIONS: 17 beta-estradiol (E2), 2'4'6'-trichloro-4-biphenylol (OH-PCB 30), 2'3'4'5'-tetrachloro-4-biphenylol (OH-PCB 61), aryl hydrocarbon receptor (AhR), cervicovaginal (CV), day of vaginal opening (DVO), diethylstilbestrol (DES), estrogen receptor (ER), hydroxylated polychlorinated biphenyls (OH-PCBs), mouse mammary tumor virus (MMTV).

Outline of manuscript section header

ABSTRACT	pg. 5
INTRODUCTION	pg. 6
MATERIALS AND METHODS	
<i>Chemicals</i>	pg. 9
<i>Animals</i>	pg. 9
<i>Statistical Analyses</i>	pg. 10
RESULTS	
<i>Gross observations</i>	pg. 11
<i>Tumor Incidence</i>	pg. 11
<i>Interactive Effects of Chemical Mixtures</i>	pg. 12
DISCUSSION	pg. 14
CONCLUSION	pg.18
REFERENCES	pg. 20
Table 1-4	pg. 28
Figure 1 legend	pg. 33
Figure 1	pg. 33

ABSTRACT

The neonatal mouse model has been a valuable tool in determining the long-term effects of early exposure to estrogenic agents in mammals. Using this model, we compared the effects of 2'4'6'-trichloro-4-biphenylol (OH-PCB 30) and 2'3'4'5'-tetrachloro-4-biphenylol (OH-PCB 61) as prototype estrogenic hydroxylated PCBs (OH-PCBs) since they are reported to exhibit relatively high estrogenic activity both *in vivo* and *in vitro*. The purpose of this study was to examine the relationship between estrogenicity and carcinogenicity of OH-PCB congeners. The OH-PCBs were tested individually and in combination to determine whether effects of combined OH-PCBs differed from those of these OH-PCBs alone. We evaluated the long-term effects of neonatal exposure to OH-PCBs with treatment doses that were based on the reported binding affinity of specific OH-PCB congeners to the estrogen receptor alpha. BALB/cCrgl female mice were treated within 16 hr after birth by subcutaneous injections every 24 hr, for 5 days. The OH-PCB 30 [200 µg/day] or 17 beta-estradiol (5 µg/day) showed similar increased incidences of cervicovaginal (CV) tract carcinomas (43 % and 47 %, respectively). In addition, when mice were treated with OH-PCBs as a mixture, a change in the type of CV tract tumor was observed, shifting from predominantly squamous cell carcinomas to adenosquamous cell carcinoma. From our results, we concluded that the individual OH-PCBs tested were estrogenic and tumorigenic in mice when exposed during development of the reproductive tract. These data support the hypothesis that mixtures may act differently and unexpectedly than as individual compounds.